

On the quantum phase problem

J.M. Vargas-Martínez and H. Moya-Cessa

¹ *INAOE, Coordinación de Óptica,*

Apdo. Postal 51 y 216, 72000 Puebla, Pue., Mexico

(Dated: February 1, 2008)

Abstract

We present a phase formalism that passes the Barnett-Pegg acid test, i.e. phase fluctuations for a number state are the expected value $\pi^2/3$ which are the fluctuations for a classical random phase distribution. The formalism is shown to have consistency subjected to different approaches.

PACS numbers: 42.50.-p, 42.65.Ky, 03.65.-w

The search for a Hermitean phase operator started with the beginning of quantum electrodynamics. The problem of phase was first addressed by Dirac [1] early in the history of quantum mechanics. However, Dirac's solution was found to suffer from mathematical difficulties [2]. Since then several formalisms have been introduced [3, 4, 5] that however have not been completely satisfactory because the way they are constructed (see Lynch [6] for a review). One of the most successful formalisms is that of Pegg and Barnett [4] that is build in a finite-dimensional Hilbert space and where all the calculations on physical quantities must be done in such a space and only after they (the calculations) have been realized the infinite-dimensional limit is taken. Not surprisingly, the Pegg-Barnett phase operator passes the Barnett-Pegg "acid test" [7] that states that the phase fluctuations for a number state must be equal to $\pi^2/3$, which is the value for a classical random phase distribution.

Other approaches based on radial integration of quasiprobability distribution functions have been proposed to describe phase properties (see for instance [8]). However, (radially) integrated distributions such as the Wigner function have shown not to work well because of its negativity [9].

Other mechanisms to describe phase have been proposed that directly write the Wigner function not in terms of position and momentum but on number and phase [10].

Here we would like to put forward a formalism for phase that passes Barnett-Pegg's acid test.

Classically we may decompose a complex c-number, A , in amplitude and phase by simply writing $A = re^{i\phi}$, with $r = |A|$ and

$$\phi = -i \ln \frac{A}{r}, \quad (1)$$

where it is implied that we have chosen the principal branch of the multi-valued logarithm function.

A Hermitean operator in correspondence to the classical form (1) was proposed by Arroyo Carrasco and Moya-Cessa [11]

$$\hat{\phi} = -\frac{i}{2} \hat{D}(\chi) \left[\ln \left(1 + \frac{\hat{a}}{\chi} \right) - \ln \left(1 + \frac{\hat{a}^\dagger}{\chi} \right) \right] \hat{D}^\dagger(\chi), \quad (2)$$

where $\hat{a}$ and $\hat{a}^\dagger$ are the annihilation and creation operator for the harmonic oscillator, respectively, $\hat{D}(\chi) = e^{\chi(\hat{a}^\dagger - \hat{a})}$ is the displacement operator with χ a real parameter to ensure convergence of the series

$$\ln \left(1 + \frac{\hat{a}}{\chi} \right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{\hat{a}}{\chi} \right)^k. \quad (3)$$

The operator (2) may be found to be Turski's operator [12] by using the unity operator given in terms of coherent states $\hat{1} = \frac{1}{\pi} \int |\alpha\rangle\langle\alpha| d^2\alpha$ and inserting this expression into (2) which yields

$$\hat{\phi} = -\frac{i}{2} \hat{D}(\chi) \left[\ln \left(1 + \frac{\hat{a}}{\chi} \right) \frac{1}{\pi} \int |\alpha\rangle\langle\alpha| d^2\alpha - \frac{1}{\pi} \int |\alpha\rangle\langle\alpha| d^2\alpha \ln \left(1 + \frac{\hat{a}^\dagger}{\chi} \right) \right] \hat{D}^\dagger(\chi), \quad (4)$$

and that may finally be written as

$$\hat{\phi} = -\frac{i}{2\pi} \int (\ln \alpha - \ln \alpha^*) |\alpha\rangle\langle\alpha| d^2\alpha. \quad (5)$$

Again, choosing the principal branch in the above equation, we can rewrite (5) as

$$\hat{\phi} = \frac{1}{\pi} \int \theta |\alpha\rangle\langle\alpha| d^2\alpha, \quad (6)$$

where $\theta = \arg(\alpha)$. It may be easily shown that the operator (6) obeys the equation of motion

$$\frac{d\hat{\phi}}{dt} = i\omega [\hat{a}^\dagger \hat{a}, \hat{\phi}] \quad (7)$$

where ω is the frequency of the harmonic oscillator. Note that for a phase operator defined in a finite dimensional Hilbert space to obey such equation of motion, the harmonic oscillator Hamiltonian should be defined also in a finite dimensional Hilbert space [13].

Dirac's original idea was to define the phase operator as the argument of the annihilation operator. We can calculate the average value of the argument of the annihilation operator, given a wave function $|\psi\rangle$, by using the Q -function defined as

$$Q(\alpha) = \frac{1}{\pi} |\langle\alpha|\psi\rangle|^2, \quad (8)$$

by recalling that

$$\langle f(\hat{a}) \rangle = \int f(\alpha) Q(\alpha) d^2\alpha \quad (9)$$

such that

$$\langle \arg(\hat{a}) \rangle = \int \arg(\alpha) Q(\alpha) d^2\alpha, \quad (10)$$

that corresponds to the average value of (6):

$$\langle \psi | \hat{\phi} | \psi \rangle = \frac{1}{\pi} \int \theta |\langle \alpha | \psi \rangle|^2 d^2 \alpha \equiv \langle \arg(\hat{a}) \rangle. \quad (11)$$

We can calculate average values of the moments of $\arg(\hat{a})$ via (9) as

$$\langle \arg^k(\hat{a}) \rangle = \int \theta^k Q(\alpha) d^2 \alpha, \quad (12)$$

such that we can compute the phase uncertainty, $\Delta\phi = \langle \arg^2(\hat{a}) \rangle - \langle \arg(\hat{a}) \rangle^2$ for a number state $|n\rangle$, yielding the result

$$\Delta\phi = \frac{\pi^2}{3}, \quad (13)$$

and where we have used

$$Q(\alpha) = \frac{1}{\pi} |\langle \alpha | n \rangle|^2 = \frac{e^{-|\alpha|^2} |\alpha|^{2n}}{\pi n!}, \quad (14)$$

i.e. giving the correct phase uncertainty expected for a state of undefined phase. We can finally write

$$\arg^k(\hat{a}) \equiv \hat{\phi}^k = \hat{\phi}^k \frac{1}{\pi} \int |\alpha\rangle \langle \alpha| d^2 \alpha \equiv \frac{1}{\pi} \int \theta^k |\alpha\rangle \langle \alpha| d^2 \alpha \quad (15)$$

and

$$e^{\hat{i}\phi} = \frac{1}{\pi} \int e^{i\theta} |\alpha\rangle \langle \alpha| d^2 \alpha \quad (16)$$

such that

$$e^{\hat{-i}\phi} e^{\hat{i}\phi} = e^{\hat{-i}\phi} \frac{1}{\pi} \int e^{i\theta} |\alpha\rangle \langle \alpha| d^2 \alpha = \frac{1}{\pi} \int e^{-i\theta} e^{i\theta} |\alpha\rangle \langle \alpha| d^2 \alpha = \hat{1} \quad (17)$$

i.e. the exponential of phase is a unitary operator within the formalism.

In conclusion, we have presented a formalism for phase that passes Barnett-Pegg's acid test giving the correct phase uncertainty for a number state.

-
- [1] P.A.M. Dirac, Proc. R. Soc. A **114**, 243 (1927).
 - [2] W.H. Louisell, Phys. Lett. **7**, 60 (1973).
 - [3] L. Susskind and J. Glogower, Physics **1**, 49 (1964).
 - [4] D.T. Pegg and S.M. Barnett, Europhys. Lett. **6**, 483 (1988); *ibid*, Phys. Rev. A **39**, 1665 (1989).
 - [5] H. Paul, Fortschr. Phys. **22**, 657 (1974).
 - [6] R. Lynch, Phys. Rep. **256** 367 (1995)

- [7] S.M. Barnett and D.T. Pegg, J. Mod. Opt. **36**, 7 (1989).
- [8] W.P. Scheleich, A. Baddilla adn H. Paul, Phys. Rev. A **45**, 6652 (1992).
- [9] B.M. Garraway and P.L. Knight, Phys. Rev. A**46**, R5346 (1992).
- [10] J.A. Vaccaro, Opt. Commun. **113** 421 (1995); Phys. Rev. A **52**, 3474 1995; see also H. Moya-Cessa, J. Optics B, in press.
- [11] M.L. Arroyo Carrasco and H. Moya-Cessa, Quant. Semiclass. Opt. **9**, L1 (1997).
- [12] L.A Turski, Physica **57** 432 (1972).
- [13] V. Buzek, a. Wilson-Gordon, P.L. Knight and W.K. Lai, Phys. Rev. A **45**, 8079 (1992).